1995 Abstract Form for Scientific Presentations SOCIETY OF MAGNETIC RESONANCE THIRD SCIENTIFIC MEETING

FUROPEAN SOCIETY FOR MAGNETIC RESONANCE IN MEDICINE AND BIOLOGY TWELFTH ANNUAL MEETING

> Nice Acropolis, NICE, FRANCE August 19-25, 1995

V	Prefer Oral Presentation but
	willing to present as a poster
	Prefer Poster but willing to mak
	Oral Presentation
	Postar Only

☐ Video required (available only for Oral Presentations)

# TOPIC CATEGORIES AUTHORS SHOULD FILL IN A NUMERICAL CATEGORY

IMAGING

# SUBCATEGORY 4

- Brain—Animal Studies Brain—Human White Matter
- Brain-Human Other
- Functional Neuro—Clinical Head, Neck, Spine and Other
- Cardiovascular
  Cardiovascular
  Cardiovascular—Clinical
  Breast/Chost

- Cenitourinary
- 12. Joints, Cartilage, Bone and Marrow 13. Vascular Imaging 14. Interventional MRI

#### SPECTROSCOPY

#### SUBCATEGORY [

- Spectroscopic Imaging Techniques
  Multinuclear Spectroscopy
  Spectroscopic Quantitation
  Human Brain—White Matter & Neuro
  Degenerative
  Human Brain—State & School & School Human Brain—Stroke & Seizure
- Human Brain—Other Animal Brain
- Cardiovascular
- Abdominal
- Addominal Musculoskeletal Tumors—Methods and Animal Models Tumors—Clinical Other (including body fluids)

## METHODOLOGY

# SUBCATEGORY [

- Angiography Flow Quantification
- Perfusion Diffusion

- Functional Neuro—Methodology and Analysis Functional Neuro—Models and Mechanisms Microscopy, Non-proton Imaging and ESR Gradients and Hardware
- RF Coils

- Rapid Imaging Spatial Localization/Imaging Motion and Artifacts
- Other Sequences/Techniques Quantitative Imaging

- 15. Quantitative imaging
  16. Image Processing
  17. 3D Display/Rendering
  18. Contrast Mechanisms/MTC
  19. Paramagnetic Contrast Agents
  20. Other Contrast Agents
- Safety/Bioeffec
- 21. Salety/Bioenects 22. Interventional MRI

#### ABSTRACT DEADLINE:

Received no later than April 11, 1995.
All copyrights to accepted abstracts become the property of the SMR. No proprietary information may be withheld by authors.

## MAIL TO:

Society of Magnetic Resonance Third Scientific Meeting 2118 Milvia Street, Suite 201 Berkeley, CA 94704, USA

Authors mailing abstracts from outside the send abstracts by express.

Please type the name and complete mailing address of the first author. Name DR. P. W. R. WOODRUFF

Address INSTITUTE OF PSYCHIATRY DE CRESPIGNY
PARK, DENMARK HILL,
PARK, DENMARK HILL, Country LONDON, SES SAF
Telephone U.K.
Fax 0717019044
Member of SMR? <b>□</b> Yes □ No
For Office Use Only:
PROGRAM #

# Effect of Selective Attention on activity within Auditory and Occipital Cortex as demonstrated using Functional MRI

P.W.R. Woodruff 1,2, R.R. Benson 1, T. Telavage 1, P.A. Bandettini 1, K.K. Kwong 1, H.C. Breiter 1, R. Howard <sup>2</sup>, J.W. Belliveau <sup>1</sup>, T. J. Brady<sup>1</sup>, B.R. Rosen<sup>1</sup>.

'Massachusetts General Hospital NMR-Center, Boston, USA & Institute of Psychiatry, London, U.K.

## Introduction

Selective attention is a process that allows some stimuli to be processed more rapidly or effectively than others. Attention is thought to operate at a series of stages in the pathway of sensory processing. Visual selective attention enhances extra-striate cortex activity demonstrable using PET (1). Electrical and magnetic recordings of human brain activity have demonstrated that attended stimuli evoke enhanced sensory responses in both visual and auditory modalities (2). Using fMRI and surface coil over the left temporal lobe, we have previously demonstrated that activation of auditory cortex was modulated by attention to auditory versus visual stimuli (3). The purpose of this study was to extend our examination to the whole head, to determine whether: 1) laterality differences in modulation of auditory cortex were observable, and 2) attention away from auditory towards visual stimuli might modulate regions in occipital cortex.

#### Methods

Four healthy right-handed male subjects 29 -38 years old were imaged whilst alternatively attending to either visual or auditory stimuli. Subjects were simultaneously presented with pseudorandom numbers from 1-9 visually and aurally and instructed to press a key when they either either heard or saw the number "8". They were cued with "listen" and "look" at transitions. A 1.5 Tesla GE Signal scanner with EPI and head coil were used with an ASE sequence (TR=1.8-2.15;170 images per slice; voxel size 3x3x5mm). Slices parallel to the Sylvian fissure were taken to include temporal and occipital lobes. Statistical maps were generated using the Kolmogorov-Smirnov statistic between conditions and threshold significance was p<0.001. Anatomical localisation was achieved using high resolution MRI images in two planes and a template derived from the Talairach Atlas for each slice.

# Results

In three of four subjects, "auditory attend" minus "visual attend" demonstrated significant areas of activation (p < 0.001) in primary auditory cortex, posterior STG and MTG. For this condition, areas of activation were greater in left than right temporal lobes for these three subjects. Signal changes ranged from 1-4%. No significant activation was noted in occipital lobes. For "visual attend" minus "auditory attend" conditions, all four subjects exhibited significant activation (p<0.001) in both occipital lobes (mainly middle occipital gyrus) with signal changes of 1-4.6% There was a tendency for

drop in signal intensity over occipital regions to coincide with transition to the "attend auditory" condition (Fig. 2).

# Conclusions

Modulation of auditory cortex activity by attention was predominantly left sided. This suggests that auditory attention shares the same laterality preference as does language. Switching attention between auditory and visual modality was accompanied by modified signal in temporal and occipital cortical regions respectively. Attention may enhance signal within the corresponding modality. Alternatively, attention within one sensory modality may reciprocally inhibit cortical brain regions subserving the other.




Fig. 1. Signal over left posterior STG during "auditory attend" minus "visual attend"




Fig. 2. Signal over left middle occipital lobe during "attend visual" minus attend auditory condition.

For Figs. 1 & 2 white bars represent baseline condition of looking at a fixation point.

# References

Heinze HJ et al., Nature 372, 543-6, 1994. Woldorff MG et al., Proc. Natl. Acad. Sci. 90, 8722-6, 1993.

Woodruff PWR et al., 1st. International Conference on Functional Mapping of the Human Brain. Paris, France, June, 1995.